

provide an operational advantage without compromising patient care. In this era of increasing fiscal constraints and improved molecular diagnostics, these data may be used to frame and inform discussions on this issue, and help guide prospective studies aimed at defining the risks and benefits of streamlining clinical care.

#### Authors' affiliations

**Rameet H Singh**, Divisions of Obstetrics and Gynecology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

**Emily J Erbeling, Jonathan M Zenilman, Khalil G Ghanem**, Divisions of Infectious Diseases, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

Funding: None.

Competing interests: None.

RHS and KGG were involved in the study design, analysis and drafting of the manuscript; EJE and JMZ were involved in the study design and revision of the manuscript.

#### REFERENCES

- 1 Rogstad KE, Ahmed-Jushuf IH, Robinson AJ, et al. MSSVD Adolescent Sexual Health Group. Standards for comprehensive sexual health services for young people under 25 years. *Int J STD AIDS* 2002;**13**:420–4.
- 2 Fenton KA, Mercer CH, Johnson AM, et al. Reported sexually transmitted disease clinic attendance and sexually transmitted infections in Britain: prevalence, risk factors, and proportionate population burden. *J Infect Dis* 2005;**191**(Suppl 1):S127–38.
- 3 Lavery S, Pugh RN, Joseph AT. The crisis in sexual health and developing genitourinary medicine services: lessons from a primary care trust. *Int J STD AIDS* 2006;**17**:37–43.
- 4 Van Leent EJM, Heijman RLJ, de Vries HJC, et al. Low time consuming STI screening for low risk STI clinic visitors. 16th Biennial Meeting of the International Society for Sexually Transmitted Diseases Research (ISSTD), July 2005, Amsterdam.
- 5 Bradbeer C, Mears A. STI services in the United Kingdom: how shall we cope? *Sex Transm Infect* 2003;**79**:435–8.
- 6 Westrom L, Joesoef R, Reynolds G, et al. Pelvic inflammatory disease and fertility. A cohort study of 1,844 women with laparoscopically verified disease and 657 control women with normal laparoscopic results. *Sex Transm Dis* 1992;**19**:185–92.
- 7 Rhoton-Vlasak A. Infections and infertility. *Prim Care Update Ob Gyns* 2000;**7**:200–6.
- 8 Buimer M, Van Doornum GJ, Ching S, et al. Detection of Chlamydia trachomatis and Neisseria gonorrhoeae by ligase chain reaction-based assays with clinical specimens from various sites: implications for diagnostic testing and screening. *J Clin Microbiol* 1996;**34**:2395–400.
- 9 Stary A, Najim B, Lee HH. Vulval swabs as alternative specimens for ligase chain reaction detection of genital chlamydial infection in women. *J Clin Microbiol* 1997;**35**:836–8.
- 10 Garland SM, Tabrizi SN. Diagnosis of sexually transmitted infections (STI) using self-collected non-invasive specimens. *Sex Health* 2004;**1**:121–6.
- 11 Garrow SC, Smith DW, Harnett GB. The diagnosis of chlamydia, gonorrhoea, and trichomonas infections by self obtained low vaginal swabs, in remote northern Australian clinical practice. *Sex Transm Infect* 2002;**78**:278–81.
- 12 Wiesenfeld HC, Lowry DL, Heine RP, et al. Self-collection of vaginal swabs for the detection of chlamydia, gonorrhea, and trichomoniasis: opportunity to encourage sexually transmitted disease testing among adolescents. *Sex Transm Dis* 2001;**28**:321–5.
- 13 Nelson DB, Bellamy S, Gray TS, et al. Self-collected versus provider-collected vaginal swabs for the diagnosis of bacterial vaginosis: an assessment of validity and reliability. *J Clin Epidemiol* 2003;**56**:862–6.
- 14 Strauss RA, Eucker B, Savitz DA, et al. Diagnosis of bacterial vaginosis from self-obtained vaginal swabs. *Infect Dis Obstet Gynecol* 2005;**13**:31–5.
- 15 Newman SB, Nelson MB, Gaydos CA, et al. Female prisoners' preferences of collection methods for testing for Chlamydia trachomatis and Neisseria gonorrhoeae infection. *Sex Transm Dis* 2003;**30**:306–19.
- 16 Amsel R, Totten PA, Spiegel CA, et al. Nonspecific vaginitis: diagnostic criteria and microbial and epidemiologic associations. *Am J Med* 1983;**74**:14–22.
- 17 Schachter J, McCormack WM, Chernesky MA, et al. Vaginal swabs are appropriate specimens for diagnosis of genital tract infection with Chlamydia trachomatis. *J Clin Microbiol* 2003;**41**:3784–9.
- 18 Smith KS, Tabrizi SN, Fethers KA, et al. Comparison of conventional testing to polymerase chain reaction in detection of Trichomonas vaginalis in indigenous women living in remote areas. *Int J STD AIDS* 2005;**16**:811–15.
- 19 Wendel KA, Erbeling EJ, Gaydos CA, et al. Trichomonas vaginalis polymerase chain reaction compared with standard diagnostic and therapeutic protocols for detection and treatment of vaginal trichomoniasis. *Clin Infect Dis* 2002;**35**:576–80.
- 20 Joeschke R, Guyatt GH, Sackett DL. Users' guides to the medical literature. III. How to use an article about a diagnostic test. B. What are the results and will they help me in caring for my patients? The Evidence-Based Medicine Working Group. *JAMA* 1994;**271**:703–7.
- 21 Simms I, Warburton F, Westrom L. Diagnosis of pelvic inflammatory disease: time for a rethink. *Sex Transm Infect* 2003;**79**:491–4.
- 22 Lee DM, Chen MY, Bradshaw CS, et al. Is routine vaginal examination necessary for asymptomatic women attending sexual health services? *Int J STD AIDS* 2006;**17**:631–2.

#### BNF for Children 2006, second annual edition

In a single resource:

- guidance on drug management of common childhood conditions
- hands-on information on prescribing, monitoring and administering medicines to children
- comprehensive guidance covering neonates to adolescents

For more information please go to [bnfc.org](http://bnfc.org)